

Anthropogenic particles in the stomach contents and liver of the freshwater fish *Squalius cephalus*

France Collard, Johnny Gasperi, Bernard Gilbert, Gauthier Eppe, Sam Azimi, Vincent Rocher, Bruno Tassin

► **To cite this version:**

France Collard, Johnny Gasperi, Bernard Gilbert, Gauthier Eppe, Sam Azimi, et al.. Anthropogenic particles in the stomach contents and liver of the freshwater fish *Squalius cephalus*. *Science of the Total Environment*, Elsevier, 2018, 643, pp.1257 - 1264. 10.1016/j.scitotenv.2018.06.313 . hal-01830268

HAL Id: hal-01830268

<https://hal-enpc.archives-ouvertes.fr/hal-01830268>

Submitted on 19 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Anthropogenic particles in the stomach contents and liver of the freshwater fish

2 *Squalius cephalus*

3
4
5
6 France Collard^{1*}, Johnny Gasperi¹, Bernard Gilbert², Gauthier Eppe², Sam Azimi³, Vincent
7 Rocher³, Bruno Tassin¹

8
9
10 ¹Laboratoire Eau Environnement et Systèmes urbains (LEESU), Université Paris-Est Créteil, Ecole
11 des ponts Paristech, AgroParisTech, 61 avenue du Général de Gaulle, 94010 Cedex Créteil, France.

12 ²Inorganic Analytical Chemistry Laboratory, B6c, Department of Chemistry, University of Liege,
13 4000 Liege, Belgium.

14 ³Syndicat Interdépartemental pour l'Assainissement de l'Agglomération Parisienne (SIAAP),
15 Direction du Développement et de la Prospective, 82 Avenue Kléber, 92700 Colombes, France.

16
17 *Corresponding author:

18 Laboratoire Eau Environnement et Systèmes Urbains (LEESU)

19 Université Paris-Est Créteil (UPEC)

20 61, Avenue du Général de Gaulle 94010 Créteil, France

21 francecollard16@gmail.com, gasperi@u-pec.fr

22 phone: + 33145171621

23 Current address: Norwegian Polar Institute (NPI), Fram Center, 9296 Tromsø, Norway

24 Abstract

25
26 Anthropogenic particles (APs) are a very broad category of particles produced directly or indirectly
27 by human activities. Their ingestion by biota is well studied in the marine environment. In contrast,
28 studies on AP ingestion in wild freshwater organisms are scarce despite high contamination levels
29 in some rivers and lakes. In this study, we aimed to evaluate the ingestion of APs and the possible
30 occurrence of APs in the liver and muscle of a freshwater fish, *Squalius cephalus*, from the Parisian
31 conurbation. After isolation, the particles were analyzed using Raman spectroscopy. In sixty
32 stomachs, eighteen APs were found, half of which were plastics and the other half were dyed
33 particles. Twenty-five percent of sampled individuals had ingested at least one AP. The mean
34 length of the APs was 2.41 mm. No significant difference was found between the sites upstream
35 and downstream of Paris. Additionally, 5% of sampled livers contained one or more APs, which
36 were characterized as microplastics (MPs). No APs were found in the muscle tissue. The majority
37 of APs isolated from stomach contents were fibers, which is similar to the findings of a previous
38 river contamination study. This highlights that fish could be more exposed to fibers than previously
39 thought and that more studies on the impacts of fiber ingestion are required. Despite their low
40 occurrence, MPs are reported, for the first time, in the liver of a wild freshwater fish species. While
41 the pathways and impacts are still unknown, MPs also occur in liver of marine mollusks and fish.
42 Physiological *in vitro* studies are needed to better evaluate the impacts of such phenomena.

43

44 Keywords

45

46 Microplastics, fibers, Seine River, European chub, muscle, liver

47

48 Introduction

49
50 Anthropogenic particles (APs) are a very broad category of particles produced directly or indirectly
51 by human activities, thus showing an anthropogenic origin. In this study, APs are defined as small
52 pieces – fragments or fibers – with an anthropogenic origin, such as plastic, dyed particles or textile
53 fibers, regardless of their size. All textile fibers have been handled by humans (e.g., dyed) and thus
54 have an anthropogenic origin, regardless of whether their basic composition is natural. All seas and
55 oceans are contaminated by APs, including plastics (Cózar et al., 2014), and so are a large number
56 of organisms (Kühn et al., 2015). Numerous studies, reviews and books (Bergmann et al., 2015)
57 report the extent of this phenomenon, particularly plastic pollution and its impact on marine
58 environments. However, although marine environments have been well studied regarding APs
59 (mainly plastics), knowledge about freshwater contamination by APs and their ecological impacts
60 remains scarce (Dris et al., 2015b). Yet, rivers continue to be the main source of microplastics
61 (MPs) (<5 mm, (Arthur et al., 2009)) to coastal ecosystems (Lima et al. 2014; Lebreton et al. 2017).

62
63 Several studies conducted in freshwater (Dris et al., 2015a; Faure et al., 2015), such as in the Seine
64 River (Gasperi et al., 2014), the River Thames (Morritt et al., 2014), the Danube River (Lechner et
65 al., 2014), the Yangtze and Hanjiang Rivers (Wang et al., 2017), the Laurentian Great Lakes
66 (Eriksen et al., 2013) and in several Chilean rivers (Rech et al., 2014), showed that freshwater is
67 contaminated, sometimes highly contaminated, by APs, especially plastics. Among others, a study
68 by Lechner et al. (2014) calculated that, in the Danube River, the global plastic mass for both MPs
69 and macroplastics was higher than the fish larvae mass. In the U.S.A., over 4 million particles are
70 estimated to be released every day from one Californian wastewater treatment plant (Mason et al.,
71 2016).

72

73 High AP concentrations could likely lead to their ingestion by aquatic organisms. Studies focusing
74 on freshwater biota contamination are very scarce. Few studies mention AP or MP ingestion by
75 wild fish species (Faure et al., 2012, 2015; Sanchez et al., 2014; Phillips and Bonner, 2015; Pazos
76 et al., 2017; Silva-Cavalcanti et al., 2017; Vendel et al., 2017) or marine mammal species
77 (Denuncio et al., 2011), and even these have focused only on stomach contents.

78 Aquatic organisms can suffer mechanical damage from ingestion of APs, e.g., obstruction or
79 starvation, as has been shown in marine mammals and birds (Beck and Barros, 1991; Pierce et al.,
80 2004; Jacobsen et al., 2010). Furthermore, aquatic organisms, including fish, can also be impacted
81 by toxicological issues including pollutant transfer (Cedervall et al., 2012; Rochman et al., 2013;
82 Wright and Kelly, 2017) or translocation of MPs into other organs such as the liver (Avio et al.,
83 2015; Collard et al., 2017a).

84

85 In this study, we analyzed the ingestion of APs by the chub *Squalius cephalus* in two highly
86 anthropized rivers, the Marne and the Seine Rivers, which both cross the Greater Paris Megacity.
87 The chub has not been studied so far in the context of AP pollution despite its omnipresence in
88 European rivers and its abundant population (Freyhof, 2014). Due to its ubiquity, the chub could
89 be a bioindicator for AP pollution in rivers. This species has a minor economic interest, but its
90 consumption by Europeans was at its maximum in 2011 (FAO, 2011). Adults are solitary, and
91 juveniles are gregarious, but both are pelagic. Generally, they feed on insects, plants and
92 crustaceans such as crayfish (Balestrieri et al., 2006; Mann, 1976; Michel and Oberdorff, 1995).
93 However, their diet varies with age and season. In the Babuna River (Republic of Macedonia), they
94 mainly feed on algae (Chrysophyceae) and insect larvae (*Ephemeroptera* and *Plecoptera*) in spring,
95 on algae (Chrysophyceae) in summer and autumn and on diatoms and other insect larvae

96 (Chironomidae) in winter (Nastova-Gjorgjioska et al., 1997). Young individuals eat primarily
97 insects (Mann, 1976), while adults generally feed on plants or crayfish and on other fish,
98 particularly in winter (Michel and Oberdorff, 1995). Moreover, the chub adapts its diet to its habitat
99 (Piria et al., 2005). For example, in Croatia, depending on the river, the diet -dominant components
100 are *Diptera* (insects), cladocerans (Djinova, 1976 cited in Piria et al., 2005), detritus (Adamek and
101 Obrdlik, 1977) or zoobenthos (Losos et al., 1980).

102
103 AP contamination of the Greater Paris Megacity rivers, e.g., the Seine and Marne Rivers, was
104 studied during a 19-month period between April 2014 and December 2015, excluding January and
105 August 2015, from upstream to downstream suburbs crossing the very dense part of the urban area
106 (Dris et al., 2018). Because of its sprawling population (12 million inhabitants), Greater Paris exerts
107 a high anthropogenic pressure on rivers and, therefore, is a site of great interest for assessing the
108 ingestion of APs by *Squalius cephalus*. The sampling was done along the continuum of the Seine
109 and Marne Rivers, allowing a comparison between the upstream and the downstream suburbs of
110 Paris. Our objectives were to evaluate the ingestion of APs, including microplastics, by European
111 chubs caught within the Parisian conurbation and to determine whether APs translocate into muscle
112 or liver tissue in this species.

113

114 Materials and methods

115


116 *Sampling*

117

118 Sixty freshwater fish of the species *Squalius cephalus* (mean length \pm SD: 296 mm \pm 62, mean
119 weight \pm SD: 340 g \pm 239) were electrofished in the Marne and Seine Rivers around Paris (Fig. 1),

120 France, between 28 August and 2 September 2016. Of the six sampling stations, three were located
 121 upstream of Paris (Gournay-sur-Marne, Maisons-Alfort and Villeneuve-Saint-Georges), and three
 122 were located downstream of Paris (Triel-sur-Seine, Levallois and Le Pecq). The three target tissues,
 123 namely, stomach (n=60), liver (n=60) and muscle (n=22), were immediately removed on board.
 124 Stomachs and livers were put in distilled water-diluted formaldehyde solution (5%), while muscles
 125 were packaged in aluminum paper and frozen at -20°C in the lab. All tissues were weighed (wet
 126 weight) before the isolation process.

127



128
 129 Figure 1. Sampling stations in the Parisian conurbation. Circles represent sampling stations and
 130 squares represent wastewater treatment plants. Number of stomach content, liver and muscle
 131 samples are shown for each station.

132

133 *Isolation of APs*

134
135 The entirety of the stomach contents and livers were used, while only a small amount of muscle
136 sample was digested (2-4 g). All tissues were degraded using sodium hypochlorite and then filtered
137 with a 5- μ m filtering membrane according to published methodology (Collard et al., 2015).
138 Nevertheless, methanol (99%), which is typically used for the centrifugation step, was replaced by
139 ethanol (99%), as methanol degrades polyvinylchloride (Collard et al., 2015). Indeed, ethanol is
140 commonly used to store biota samples or microplastics once analyzed (Sanchez et al., 2014;
141 Phillips and Bonner, 2015; Ory et al., 2017).

142 Consequently, in the frame of this study, tissues were put into a 14 g/L NaClO solution overnight.
143 The NaClO solution was filtered with a cellulose acetate filter membrane (5- μ m porosity), which
144 was then rinsed with ethanol. This solution, containing APs, was centrifuged at 5,000 rpm for 10
145 min. The bottom was then collected and deposited onto a stainless steel plate for Raman
146 spectroscopy analysis.

147
148 *Raman analysis*

149
150 All found particles were analyzed using a LabRam 300 spectrometer (Jobin-Yvon) equipped with
151 an Olympus confocal microscope and an Andor BRDD Du401 CCD detector. Depending on the
152 color of the particle, either a Spectraphysics argon ion laser (514.5 nm) or a Torsana diode laser
153 (784.7 nm), with two different objectives (magnification of $\times 50$ or $\times 100$), was used. The maximum
154 beam laser powers induced on the sample were 5 mW (green laser) and 30 mW (red laser), but
155 several neutral density filters were used to decrease the power and avoid degradation of the sample.
156 The integration times ranged from 5 s to 50 s, depending on the sample. Two spectral databases

157 were used to perform matchings: a commercially available database (Omic Spectra software,
158 Thermo Fisher Scientific, U.S.A.) and a personal library, which used the Thermo Spectra 2.0
159 software.

160 After Raman analyses, APs were isolated in 1 ml of 99% ethanol for further observations and
161 measurements. Only particles whose anthropogenic origin was confirmed by Raman spectroscopy
162 (plastic particles, dyed particles, etc.) were included in the results. Those particles showing an
163 anthropogenic origin have been classified as “APs”. Inside the “AP” category, we defined two
164 subcategories: microplastics and other APs (e.g., dyed particles of an unknown material). In
165 addition, we classified all APs based on their shape: either fragments or fibers.

166

167 *Preventing contamination and procedural blanks*

168

169 To minimize contamination, white cotton lab coats and latex gloves were worn throughout the
170 entire isolation process. All work surfaces and dissection materials were rinsed with distilled water,
171 and all glassware was rinsed with distilled water before each use. Stainless steel plates were placed
172 under a metal sifter with 36- μ m mesh to prevent airborne contamination while the plates dried. No
173 particles smaller than 36 μ m were found. The NaClO was filtered with a 5- μ m cellulose acetate
174 membrane before dilution with distilled water.

175 Three 50-ml volumes of formaldehyde used to store the samples were filtered with the same filter
176 membrane as that used for the isolation process. No APs were found. In addition, guts and livers
177 were kept closed in the formaldehyde but were opened under an airflow cabinet just before the
178 isolation process. Muscles were kept in aluminum, but particles from this material were not found
179 in any samples.

180 Three procedural blanks were performed, along with runs of several samples, meaning that the
181 entire isolation process was verified. It began in the formaldehyde solution used to store the
182 samples and ended with the observation of stainless steel plates using Raman spectroscopy. This
183 allowed us to evaluate whether solutions, materials or equipment were contaminated by APs in any
184 way. Two particles made of an unknown organic material and two others made of only cellulose
185 were found, meaning that those particles were not APs.

186

187 *Images and measurements of particles*

188

189 APs were collected from the Raman plates with 99% ethanol solution and then put into a microtube
190 for further filtration using the same white filter membranes as those used for the isolation process.
191 APs were photographed on both the stainless steel plate and on the filter membrane using a
192 stereomicroscope (Leica MZ12, Leica AG Camera, Germany). They were then measured at their
193 longest dimension using the Histolab software (Histolab Products AB, Sweden).

194

195 *Statistics*

196

197 Statistical analyses were performed with the GraphPad Prism software (v5.03, GraphPad software
198 Inc., California, U.S.A.). Pearson's test was applied to evaluate the correlation between fish and
199 AP lengths. Normality of the data was checked using the Kolmogorov-Smirnov test prior to
200 analysis. Fisher's test was applied to compare (1) the number of fish that ingested (or not) at least
201 one AP, MP, fiber or fragment, with each type of particle considered independently; (2) the number
202 of all ingested APs versus MPs; (3) the number of ingested fibers versus fragments; and (4) the
203 concentrations of all APs and MPs in SC, between upstream and downstream of Paris. A Mann-

204 Whitney U test was applied to compare the lengths of ingested APs between the upstream and
205 downstream suburbs of Paris. All alpha values were set at 0.05. Data are expressed in mean \pm
206 standard deviation (SD).

207

208 Results

209


210 *Stomach contents (SCs)*

211

212 All isolated particles were analyzed, and 18 APs out of 70 isolated particles were found (Fig. 2).
213 The fifty-two other particles were made of natural organic polymers such as cellulose or could not
214 be identified. Twenty-five percent of fish have ingested at least one AP, and 15% have ingested at
215 least one plastic particle.

216

217



218
 219 Figure 2. Pictures of APs found in stomach contents (A-D) and livers (E, F) of *S. cephalus*. A: blue
 220 dyed fiber, B: polypropylene (PP) fiber, C: polyethylene terephthalate fiber, D: PP fiber, E:
 221 polystyrene fragment, F: polyethylene fragment. Scale bar: 500 μm .

222

223 Table 1. Characteristics of the 18 APs found in stomach contents (SC). *Concerns only fibers.

Station	Composition	Size (mm)	Diameter* (μm)	Shape	Item/g SC
Gournay-sur-Marne	PET	2.96	11	FI	0.38
	Black dyed fiber	0.84	22	FI	2.86
	PAN	7.38	30	FI	
Maisons-Alfort	Blue dyed fragment	2.41	-	FR	0.23


	Blue dyed fiber	0.96	18	FI	
	PET	0.74	12	FI	0.71
Villeneuve-Saint-Georges	PET	2.51	40	FI	2.5
	PP	0.69	59	FI	0.38
	PET	0.39	32	FI	0.67
	Blue dyed fiber	1.08	27	FI	0.38
Triel-sur-Seine	Blue dyed particle	1.43	23	FI	1.7
	Blue dyed fiber	2.21	17	FI	
	PEVA	6.56	-	FR	0.36
	PP	0.44	-	FR	0.67
Levallois	PET	2.36	56	FI	1.0
	Blue dyed fiber	4.44	22	FI	0.59
Le Pecq	Blue dyed fiber	1.21	31	FI	3.33
	Black dyed fiber	4.79	22	FI	2.0

224 PET=polyethylene terephthalate, PP=polypropylene, PEVA=poly(ethylene-co-vinyl acetate),

225 PAN=polyacrylonitrile, FI=fiber, FR=fragment.

226


227



228
 229 Figure 3. Summary of the main characteristics of APs found in stomach contents. a: total AP
 230 amount according to shape, b: total AP amount according to composition.

231
 232 Of the 18 APs, 15 were fibers, and 3 were fragments (Fig. 3a). Fibers longer than 1 mm were
 233 dominant (n=10), followed by fibers ranging from 0.5 to 1 mm (n=4) and by fibers smaller than
 234 0.5 mm (n=1). The mean diameter of fibers was $28 \pm 14 \mu\text{m}$. Among the 9 plastic particles, 7 were
 235 fibers, and two were fragments. APs measured on average $2.41 \pm 2.09 \text{ mm}$, with a minimum length
 236 of $390 \mu\text{m}$ and a maximum length of 7.38 mm. Nine APs were made of plastic polymers and nine
 237 were other kinds of APs, i.e. dyed particles (Table 1, Fig. 3b). Four types of plastic polymers were
 238 found (Fig. 4): polyethylene terephthalate (PET, n=5), polypropylene (PP, n=2), polyacrylonitrile
 239 (PAN, n=1) and poly(ethylene-co-vinyl acetate) (PEVA, n=1). Furthermore, twenty fibers of
 240 cellulose were found in SC samples but were not included in APs, because the anthropogenic origin
 241 could not be confirmed using Raman spectroscopy. There is no correlation between the length of
 242 ingested APs and the length of fish ($R^2=0.0026$).

243
 244



245
 246 Figure 4. Raman spectra obtained from isolated APs. PET=polyethylene terephthalate,
 247 PP=polypropylene, PEVA=poly(ethylene-co-vinyl acetate), PAN=polyacrylonitrile.

248
 249 When comparing the areas upstream and downstream of Paris, no significant difference was found
 250 regarding the number of individuals having ingested MPs alone, all APs, fibers and fragments; the
 251 number of ingested APs and MP; the number of ingested fibers and fragments (contingency tables,
 252 $\alpha > 0.05$); and the length of all ingested APs (Mann-Whitney U test, $\alpha = 0.24$). A summary of the
 253 results is shown in Table 2. Fish from the upstream stations had 0.32 AP/g of SC, while fish from
 254 the downstream stations had 0.40 AP/g of SC, also showing no significant difference (contingency
 255 tables, $\alpha > 0.05$).

256
 257 Table 2. Comparison of different data between the areas upstream and downstream of Paris.
 258 %=percentage.

	Upstream (n=33)	Downstream (n=27)
% of fish with AP	24%	26%
% of fish with MP	18%	11%
Mean AP size	2.00 mm	2.93 mm
Min-max size	0.39-7.38 mm	0.44-6.56 mm
% of fibers	90%	75%
% of fragments	10%	15%
AP/g of SC (wet weight)	0.32	0.40

259

260

261 *Livers & Muscles*

262

263 Four APs, of 13 total extracted particles, were found in three livers out of the sixty analyzed. All
 264 were MPs, but three were made of PE, and one was made of polystyrene (PS). Their lengths ranged
 265 from 147 μm to 567 μm (Table 3). Given the size of these MP fragments, it is very unlikely that
 266 they come from airborne contamination.

267

268 Table 3. Characteristics of MPs found in livers.

Station	Polymer	Size (mm)	Shape
Levallois	PE	0.567	FR
	PE	0.161	FR
	PE	0.147	FR

Villeneuve- Saint- Georges	PS	0.270	FR
---	----	-------	----

269 FR=fragment, PE=polyethylene, PS=polystyrene.

270

271 No APs were found in muscle tissue.

272

273 Discussion

274

275 No difference was found between the areas upstream and downstream of Paris. Regarding shape,
 276 fibers were dominant, and fibers longer than 1 mm were found in a higher quantity than the two
 277 other size classes. The smallest fiber found in SCs was 390 μm long. Smaller particles are unlikely
 278 to be ingested because they would probably be rejected with water through branchial structures
 279 (Collard et al., 2017b). Although Dris et al. (2015a) did not use any spectroscopic method for
 280 identification, these three results are consistent with those highlighted by the previous study in the
 281 Seine River water. Their study found an average of 45 fibers/ m^3 and 0.54 fragments/ m^3 in the water
 282 column but found no difference between the downstream and upstream areas, and fibers longer
 283 than 1 mm were predominant. European chubs' stomach contents seem to reflect what is found in
 284 the river water. Moreover, since it is not an endangered species in European waters, European chub
 285 could then be used as a bioindicator for AP and MP pollution. It is already used as a common
 286 bioindicator in European freshwaters for other pollutants, such as musk (Hájková et al., 2007),
 287 metals (Dragun et al., 2016), dioxins and polychlorobiphenyls (Pacini et al., 2013).

288

289 Compared to fibers, only a few anthropogenic fragments were found. Fragments may be indirectly
290 ingested through various pathways: transfer from prey, as shown in experimental studies (Cedervall
291 et al., 2012; Farrell and Nelson, 2013; Setälä et al., 2014); accidental ingestion from the water
292 column or from sediments; or intentional ingestion (Ory et al., 2017). The chub is an omnivorous
293 (Piria et al., 2005) and opportunistic fish (Balestrieri et al., 2006), meaning that everything is
294 potential prey to feed on, including plastic. However, mechanisms of ingestion cannot be elucidated
295 in this study.

296
297 All former studies dealing with AP ingestion by freshwater fish focused on MPs. In comparison,
298 the next section will also focus on MP ingestion but not on all APs. Few studies exist on MP
299 ingestion by freshwater fishes, and moreover, their results must be carefully compared. Locations,
300 species, and methodologies, including characterization of the target particles, differ between
301 studies, leading to inconsistent comparisons. Here, we chose to provide data in terms of
302 items/individual, items/g SC and percentage of occurrence. Each of those units has advantages and
303 disadvantages but will at least allow comparisons between previous and future studies. The
304 percentage of MP occurrence is often used to make comparisons more reliable. Regarding MP
305 ingestion by freshwater fish, some of the lowest percentages of MP occurrence are found in fish
306 from Geneva Lake (Faure et al., 2015), from French rivers (Sanchez et al., 2014), from a Brazilian
307 estuary (Vendel et al., 2017), and in European smelts *Osmerus eperlanus* from the Thames estuary
308 (McGoran et al., 2017) (7.5%, 12%, 9%, and 20% of individuals had ingested plastics,
309 respectively). With an MP occurrence rate of 15%, our study has one of the lowest percentages. In
310 addition, some studies have reported high MP occurrence percentages of up to 100% (Pazos et al.,
311 2017). High values were also found in the Thames estuary, where 71% of flounders *Platichthys*
312 *flesus* have ingested MPs (McGoran et al., 2017). Flounders are a benthic fish, raising questions

313 about the possible difference between pelagic and benthic feeders. According to a study on
314 European smelts (pelagic) and flounders (benthic), feeding strategies could have an influence on
315 the ingestion of APs (McGoran et al., 2017). The benthic species had a much higher ingestion rate
316 than the pelagic one, but this was not statistically assessed. By contrast, Jabeen et al. (2017) and
317 Murphy et al. (2017) found a significant difference between demersal and pelagic fishes (in
318 freshwaters and marine waters, respectively). The demersal species ingested more APs. However,
319 our study concerned a pelagic species, and we found a relatively low rate of MP occurrence
320 compared to high percentages previously cited. Our percentage of MP occurrence in SC could have
321 been underestimated. Among the dyed particles found in our study, some may have been made of
322 a plastic polymer, which cannot be detected using Raman spectroscopy because of the dye coating.
323 In marine environments, sediments are a sink for microplastics (Woodall et al., 2014). Regarding
324 freshwaters, the literature lacks information about river sediment microplastic contamination,
325 highlighting the need for more studies to assess whether freshwater demersal feeders are more
326 exposed to pelagic ones.

327
328 In the SC, a higher proportion of fibers was found compared to fragments (83% vs. 17%), as
329 expected based on other studies (McGoran et al., 2017; Pazos et al., 2017; Silva-Cavalcanti et al.,
330 2017; Vendel et al., 2017). As the digestion protocol may have bleached some particles because of
331 the use of NaClO, some dyed cellulosic fibers may have, in actuality, been dyed textile fibers that
332 were bleached during the isolation process, leading to an underestimation of APs. However, the
333 results concerning the type of particle ingested by chubs are consistent with what is commonly
334 found in the Seine and Marne Rivers: fibers are much more abundant than fragments.

335 WWTPs could be sources of fibers in urban rivers. While they help to retain a proportion of them,
336 e.g., from 83% to 95% (Dris et al. 2015a), WWTP effluents still constitute as a source of fibers

337 (Dris et al. 2015a; Leslie et al. 2017). Paris' WWTPs remove long fibers (> 1 mm), but smaller
338 fibers can pass through (Dris et al., 2015a). Atmospheric fallout can also be an additional source
339 of fibers in receiving systems. Indeed, Dris et al. (2016) found that 50% of fibers coming from
340 atmospheric fallout were longer than 1 mm, which is the dominant size class ingested by chubs. In
341 that study, 50% of all fibers were natural, 21% were manufactured by transformation of natural
342 polymers (e.g., rayon), and 29% contain a petrochemical-derived polymer (e.g., mix of cotton and
343 polyamide, or polyurethane). No information regarding the >1 mm size class was given. Even if
344 no direct link can be established between atmospheric fallout and the presence of long fibers in
345 chubs' SC, we speculate that atmospheric inputs in water could be an important source of the
346 longest fibers, which could then be ingested by freshwater biota.

347 The release of fibers into the environment is already a major concern, since washing machines can
348 release a high amount of fibers per wash (Hartline et al., 2016; Hernandez et al., 2017). A polyester
349 garment can release fibers between 0.033% and 0.039% of its weight (Dubai and Liebezeit,
350 2013), and six identical fleece garments can release fibers from 0.008% to 0.021% of their total
351 weight (Pirc et al., 2016). Consequently, in urban environments, fibers may be more problematic
352 than fragments, particularly for filtering organisms that have adapted their morphology to retain
353 such particles (Collard et al., 2017b).

354
355 An increasing number of studies report AP, mainly MP, ingestion by fish, but the impacts are
356 unknown. Studies which focus on wild-caught species probably have data regarding only a
357 snapshot of the fish's lifetime. After being ingested, MPs are probably excreted (Van
358 Cauwenberghe and Janssen, 2014; Watts et al., 2014; Grigorakis et al., 2017), but they may have
359 a longer retention time than food that is typically (Ward and Kach, 2009; Mazurais et al., 2015).
360 Additionally, no mechanical harm resulting from MPs has been reported in fish, but in birds,

361 plastics can reduce hunger or cause a reduction in assimilation efficiency (Ryan, 1989; Lenzi et al.,
362 2016). The problem may instead concern pollutants. Microplastic fibers, like other microplastics,
363 adsorb pollutants present in the surrounding water and contain small amounts of additives. They
364 can be transferred to organisms once ingested (Teuten et al., 2009), although this is controversial
365 (Koelmans et al., 2016). Additionally, little is known about adsorption on APs other than those that
366 are plastic. In our study, 9 out of 18 APs were not made of plastic. How do pollutants interact with
367 such particles? Did they also leach from particles to organs? Fibers constitute a major part of AP
368 pollution in freshwater environments, so answering these questions is a priority for AP pollution
369 research.

370
371 For the first time, MPs have been found in the livers of wild freshwater fish. Translocation also
372 occurs in Parisian freshwaters, where fragment contamination is lower than fiber contamination
373 (Dris, 2016). Translocation phenomena of such large MP abundances in fish have already occurred
374 in both the laboratory (Avio et al., 2015) and the marine environment (Collard et al., 2017a). Other
375 organisms also exhibit MP translocation (von Moos et al., 2012; Brennecke et al., 2015) but in the
376 laboratory only. In mammals, the uptake of microparticles from the gut and subsequent passage
377 through Peyer's patches in the intestine mucosa has been known for several decades (LeFevre et
378 al., 1978; Jani et al., 1989; Jani et al., 1990). However, fish do not contain Peyer's patches,
379 (Rombout et al., 1993; Cairn and Swan, 2010), and MP translocation pathways through the
380 intestinal barrier are unknown in these organisms. Some hypotheses have been proposed, such as
381 persorption, during which particles pass between cells of the gut epithelium to the circulatory
382 system (Volkheimer, 1975; Wright and Kelly, 2017); endocytosis, which is an internalization of
383 particles in cells of the digestive gland and has been demonstrated in mussels (Von Moos et al.,
384 2012); and, finally, specialized enterocytes called microfold cells may be involved, as suggested

385 by Browne et al. (2008), again, in mussels. The digestive system in mussels differs significantly
386 from that of fish, but since the literature regarding MP translocation in fish is very scarce, we can
387 only rely on those few studies to piece together an explanation. Considering the large sizes of MPs
388 found in livers in this study, the first hypothesis could be the most relevant. Persorption has been
389 observed for particles of up to 130 μm (Steffens, 1995 cited in Wright and Kelly, 2017). The
390 remaining possibilities seem unlikely considering the sizes of translocated particles: 3 μm (Browne
391 et al., 2008) and smaller than 80 μm (Von Moos et al., 2012). However, this should be confirmed
392 by experimental studies.

393
394 No APs were detected in muscle tissue despite the use of the sensitive Raman-based method
395 developed. However, fewer samples, compared to those of SC and liver, were collected, and only
396 several grams of muscle tissue per individual were taken to limit the amount of solution used.
397 Consequently, other studies focusing on contamination of muscle tissue, and of other organs, are
398 needed.

399
400 Conclusions

401
402 In conclusion, the opportunistic European chub ingested APs in an amount similar to that observed
403 in other freshwater species (McGoran et al., 2017). Our results reflected the main ones highlighted
404 in a study performed in the Seine and Marne Rivers inhabited by chubs. We then suggested that
405 chubs may be used as an indicator species, as they are not endangered but are in fact common
406 throughout Europe. Half of the analyzed APs were found to be made of plastic, highlighting that
407 focusing on plastic particles could underestimate the level of APs ingested, regardless of species.
408 Fibers were the predominant shape found in stomach contents and are ingested with unknown

409 consequences. Fragments can translocate into other organs by unknown pathways, as shown in
410 other studies. Until now, this was not observed for fibers.

411 Many questions are yet to be answered. In the field, do MPs translocate into organs other than the
412 liver? What are the pathways of translocation? How do they impact the organism? Does the MP
413 contamination level in the environment influence the translocation process? What is the proportion
414 of ingested plastic that can translocate into the liver? Considering the current global consumption
415 of fish, humans are directly impacted by those issues. Histological analyses could provide answers
416 to these questions. Histological analyses could contribute to give answers to these questions. The
417 next step in AP contamination research could be to focus on *in vitro* studies, which may help to
418 evaluate impacts and determine what type(s) of particles are likely to be translocated and the
419 reasons for such translocation (polymer, size, presence of dyes on surface molecules, etc.).

420

421 Acknowledgments

422

423 The research leading to these results received funding from the People Programme (Marie Curie
424 Actions) of the European Union's Seventh Framework Programme (FP7/2007-2013) under the
425 REA grant agreement n. PCOFUND-GA-2013-609102, through the PRESTIGE programme
426 coordinated by Campus France (postdoctoral grant). The authors also want to thank the University
427 of Paris-Est Créteil (UPEC) and the SIAAP scientists who helped during the sampling survey.

428

429

430 References

- 431
- 432 Adamek, Z., Obrdlik, P., 1977. Food of important cyprinid species in the warmed barb zone of
433 the Oslava river. *Folia Zool.* 26, 171–182.
- 434 Arthur, C., Baker, J., Bamford, H. (eds), 2009. Proceedings of the International Research
435 Workshop on the Occurrence, Effects and Fate of Microplastics Marine Debris.
- 436 Avio, C.G., Gorbi, S., Regoli, F., 2015. Experimental development of a new protocol for
437 extraction and characterization of microplastics in fish tissues: First observations in
438 commercial species from Adriatic Sea. *Mar. Environ. Res.* 111, 18–26.
439 <https://doi.org/10.1016/j.marenvres.2015.06.014>
- 440 Balestrieri, A., Prigioni, C., Remonti, L., Sgrosso, S., Priore, G., 2006. Feeding ecology of
441 *Leuciscus cephalus* and *Rutilus rubilio* in southern Italy. *Ital. J. Zool.* 73, 129–135.
442 <https://doi.org/10.1080/11250000600679561>
- 443 Barreiros, J.P., Raykov, V.S., 2014. Lethal lesions and amputation caused by plastic debris and
444 fishing gear on the loggerhead turtle *Caretta caretta* (Linnaeus, 1758). Three case reports
445 from Terceira Island, Azores (NE Atlantic). *Mar. Pollut. Bull.* 86, 518–22.
446 <https://doi.org/10.1016/j.marpolbul.2014.07.020>
- 447 Beck, C.A., Barros, N.B., 1991. The impact of debris on the Florida manatee. *Mar. Pollut. Bull.*
448 22, 508–510. [https://doi.org/http://dx.doi.org/10.1016/0025-326X\(91\)90406-I](https://doi.org/http://dx.doi.org/10.1016/0025-326X(91)90406-I)
- 449 Bergmann, M., Gutow, L., Klages, M., 2015. Marine anthropogenic litter. Springer.
- 450 Brennecke, D., Ferreira, E.C., Costa, T.M.M., Appel, D., da Gama, B.A.P., Lenz, M., 2015.
451 Ingested microplastics (>100µm) are translocated to organs of the tropical fiddler crab
452 *Uca rapax*. *Mar. Pollut. Bull.* 96, 491–495. <https://doi.org/10.1016/j.marpolbul.2015.05.001>
- 453 Browne, M.A., Dissanayake, A., Galloway, T.S., Lowe, D.M., Thompson, R.C., 2008. Ingested

454 Microscopic Plastic Translocates to the Circulatory System of the Mussel, *Mytilus edulis*
455 (L.). Environ. Sci. Technol. 42, 5026–5031. <https://doi.org/10.1021/es800249a>

456 Cairn, K., Swan, C., 2010. Fish Physiology: The Multifunctional Gut of Fish, Fish Physiology.
457 Elsevier Science.

458 Cedervall, T., Hansson, L.-A., Lard, M., Frohm, B., Linse, S., 2012. Food chain transport of
459 nanoparticles affects behaviour and fat metabolism in fish. PLoS One 7, e32254.
460 <https://doi.org/10.1371/journal.pone.0032254>

461 Collard, F., Gilbert, B., Compère, P., Eppe, G., Das, K., Jauniaux, T., Parmentier, E., 2017a.
462 Microplastics in livers of European anchovies (*Engraulis encrasicolus*, L.). Environ. Pollut.
463 229, 1000–1005. <https://doi.org/10.1016/j.envpol.2017.07.089>

464 Collard, F., Gilbert, B., Eppe, G., Parmentier, E., Das, K., 2015. Detection of Anthropogenic
465 Particles in Fish Stomachs: An Isolation Method Adapted to Identification by Raman
466 Spectroscopy. Arch. Environ. Contam. Toxicol. 69, 331–339.
467 <https://doi.org/10.1007/s00244-015-0221-0>

468 Collard, F., Gilbert, B., Eppe, G., Roos, L., Compère, P., Das, K., Parmentier, E., 2017b.
469 Morphology of the filtration apparatus of three planktivorous fishes and relation with
470 ingested anthropogenic particles. Mar. Pollut. Bull. 116, 182–191.
471 <https://doi.org/10.1016/j.marpolbul.2016.12.067>

472 Cózar, A., Echevarría, F., González-Gordillo, J.I., Irigoien, X., Ubeda, B., Hernández-León, S.,
473 Palma, A.T., Navarro, S., García-de-Lomas, J., Ruiz, A., Fernández-de-Puelles, M.L.,
474 Duarte, C.M., 2014. Plastic debris in the open ocean. Proc. Natl. Acad. Sci. U. S. A. 111,
475 10239–10244. <https://doi.org/10.1073/pnas.1314705111>

476 Denuncio, P., Bastida, R., Dassis, M., Giardino, G., Gerpe, M., Rodríguez, D., 2011. Plastic
477 ingestion in Franciscana dolphins, *Pontoporia blainvillei* (Gervais and d’Orbigny, 1844),

478 from Argentina. Mar. Pollut. Bull. 62, 1836–1841.
479 <https://doi.org/10.1016/j.marpolbul.2011.05.003>

480 Djinova, L., 1976. Za ishranata za klenot (*Leuciscus cephalus macedonicus* Kar.) i mrenata
481 (*Barbus cyclolepis strumicae* Kar.) od rekata i od akumulacijata Turija. Folia Balc. 3.

482 Dragun, Z., Tepić, N., Krasnići, N., Teskeredžić, E., 2016. Accumulation of metals relevant for
483 agricultural contamination in gills of European chub (*Squalius cephalus*). Environ. Sci.
484 Pollut. Res. 23, 16802–16815. <https://doi.org/10.1007/s11356-016-6830-y>

485 Dris, R., 2016. First assessment of sources and fate of macro- and micro- plastics in urban
486 hydrosystems: Case of Paris megacity. Université de Paris-Est Créteil.

487 Dris, R., Gasperi, J., Saad, M., Mirande, C., Tassin, B., 2016. Synthetic fibers in atmospheric
488 fallout: A source of microplastics in the environment? Mar. Pollut. Bull. 104, 290–293.
489 <https://doi.org/10.1016/j.marpolbul.2016.01.006>

490 Dris, R., Gasperi, J., Rocher, V., Saad, M., Renault, N., Tassin, B., 2015a. Microplastic
491 contamination in an urban area: A case study in Greater Paris. Environ. Chem. 12, 592–599.
492 <https://doi.org/10.1071/EN14167>

493 Dris, R., Gasperi, J., Rocher, V., Tassin, B., 2018. Synthetic and non-synthetic anthropogenic
494 fibers in a river under the impact of Paris Megacity: Sampling methodological aspects and
495 flux estimations. Sci. Total Environ. 618, 157–164.
496 <https://doi.org/10.1016/j.scitotenv.2017.11.009>

497 Dris, R., Imhof, H., Sanchez, W., Gasperi, J., Galgani, F., Tassin, B., Laforsch, C., 2015b.
498 Beyond the ocean: Contamination of freshwater ecosystems with (micro-)plastic particles.
499 Environ. Chem. 12, 539–550. <https://doi.org/10.1071/EN14172>

500 Dubaish, F., Liebezeit, G., 2013. Suspended microplastics and black carbon particles in the Jade
501 system, southern North Sea. Water. Air. Soil Pollut. 224.

502 Eriksen, M., Mason, S., Wilson, S., Box, C., Zellers, A., Edwards, W., Farley, H., Amato, S.,
503 2013. Microplastic pollution in the surface waters of the Laurentian Great Lakes. *Mar.*
504 *Pollut. Bull.* 77, 177–182. <https://doi.org/10.1016/j.marpolbul.2013.10.007>

505 FAO, 2011. FAO Capture Production of *Leuciscus Cephalus* [WWW Document]. URL
506 <http://www.fishbase.se/report/FAO/FAOCatchList.php?scientific=Leuciscus+cephalus>
507 (accessed 9.15.17).

508 Farrell, P., Nelson, K., 2013. Trophic level transfer of microplastic: *Mytilus edulis* (L.) to
509 *Carcinus maenas* (L.). *Environ. Pollut.* 177, 1–3.
510 <https://doi.org/10.1016/j.envpol.2013.01.046>

511 Faure, F., Corbaz, M., Baecher, H., de Alencastro, L., 2012. Pollution due to plastics and
512 microplastics in Lake Geneva and in the Mediterranean Sea. *Arch. Sci.* 65, 157–164.

513 Faure, F., Demars, C., Wieser, O., Kunz, M., De Alencastro, L.F., 2015. Plastic pollution in
514 Swiss surface waters: Nature and concentrations, interaction with pollutants. *Environ.*
515 *Chem.* 12, 582–591. <https://doi.org/10.1071/EN14218>

516 Freyhof, J., 2014. The IUCN Red List of Threatened Species 2014: *Squalius cephalus* [WWW
517 Document]. URL <http://www.iucnredlist.org/details/61205/0> (accessed 9.15.17).

518 Gasperi, J., Dris, R., Bonin, T., Rocher, V., Tassin, B., 2014. Assessment of floating plastic
519 debris in surface water along the Seine River. *Environ. Pollut.* 195, 163–166.
520 <https://doi.org/10.1016/j.envpol.2014.09.001>

521 Grigorakis, S., Mason, S.A., Drouillard, K.G., 2017. Determination of the gut retention of plastic
522 microbeads and microfibers in goldfish (*Carassius auratus*). *Chemosphere* 169, 233–238.
523 <https://doi.org/http://dx.doi.org/10.1016/j.chemosphere.2016.11.055>

524 Hájková, K., Pulkrabová, J., Hajšlová, J., Randák, T., Žlábek, V., 2007. Chub (*Leuciscus*
525 *cephalus*) as a Bioindicator of Contamination of the Vltava River by Synthetic Musk

526 Fragrances. *Arch. Environ. Contam. Toxicol.* 53, 390–396. <https://doi.org/10.1007/s00244->
527 006-0190-4

528 Hartline, N.L., Bruce, N.J., Karba, S.N., Ruff, E.O., Sonar, S.U., Holden, P.A., 2016. Microfiber
529 Masses Recovered from Conventional Machine Washing of New or Aged Garments.
530 *Environ. Sci. Technol.* 50, 11532–11538. <https://doi.org/10.1021/acs.est.6b03045>

531 Hernandez, E., Nowack, B., Mitrano, D.M., 2017. Polyester Textiles as a Source of Microplastics
532 from Households: A Mechanistic Study to Understand Microfiber Release During Washing.
533 *Environ. Sci. Technol.* 51, 7036–7046. <https://doi.org/10.1021/acs.est.7b01750>

534 Jabeen, K., Su, L., Li, J., Yang, D., Tong, C., Mu, J., Shi, H., 2017. Microplastics and
535 mesoplastics in fish from coastal and fresh waters of China. *Environ. Pollut.* 221, 141–149.
536 <https://doi.org/http://dx.doi.org/10.1016/j.envpol.2016.11.055>

537 Jacobsen, J.K., Massey, L., Gulland, F., 2010. Fatal ingestion of floating net debris by two sperm
538 whales (*Physeter macrocephalus*). *Mar. Pollut. Bull.* 60, 765–7.
539 <https://doi.org/10.1016/j.marpolbul.2010.03.008>

540 Jani, P., Halbert, G.W., Langridge, J., Florence, A.T., 1990. Nanoparticle uptake by the rat
541 gastrointestinal mucosa: Quantitation and particle size dependency. *J. Pharm. Pharmacol.*
542 42, 821–826.

543 Jani, P., Halbert, G.W.W., Langridge, J., Florence, A.T.T., 1989. The Uptake and Translocation
544 of Latex Nanospheres and Microspheres after Oral Administration to Rats. *J. Pharm.*
545 *Pharmacol.* 41, 809–812. <https://doi.org/10.1111/j.2042-7158.1989.tb06377.x>

546 Koelmans, A.A., Bakir, A., Burton, G.A., Janssen, C.R., 2016. Microplastic as a Vector for
547 Chemicals in the Aquatic Environment: Critical Review and Model-Supported
548 Reinterpretation of Empirical Studies. *Environ. Sci. Technol.* 50, 3315–3326.
549 <https://doi.org/10.1021/acs.est.5b06069>

550 Kühn, S., Bravo Rebolledo, E.L., Van Franeker, J.A., 2015. Deleterious Effects of Litter on
551 Marine Life, in: Bergmann, M., Gutow, L., Klages, M. (Eds.), *Marine Anthropogenic Litter*.
552 Berlin, p. 447. https://doi.org/http://dx.doi.org/10.1007/978-3-319-16510-3_4

553 Lebreton, L.C.M., van der Zwet, J., Damsteeg, J.-W., Slat, B., Andrady, A., Reisser, J., 2017.
554 River plastic emissions to the world's oceans. *Nat. Commun.* 8, 15611.
555 <https://doi.org/10.1038/ncomms15611>

556 Lechner, A., Keckeis, H., Lumesberger-Loisl, F., Zens, B., Krusch, R., Tritthart, M., Glas, M.,
557 Schludermann, E., 2014. The Danube so colourful: A potpourri of plastic litter outnumbers
558 fish larvae in Europe's second largest river. *Environ. Pollut.* 188, 177–181.
559 <https://doi.org/http://dx.doi.org/10.1016/j.envpol.2014.02.006>

560 LeFevre, M.E., Vanderhoff, J.W., Laissue, J.A., Joel, D.D., 1978. Accumulation of 2- μ m latex
561 particles in mouse Peyer's patches during chronic latex feeding. *Experientia* 34, 120–122.
562 <https://doi.org/10.1007/BF01921939>

563 Lenzi, J., Burgues, M.F., Carrizo, D., Machín, E., Teixeira-de Mello, F., 2016. Plastic ingestion
564 by a generalist seabird on the coast of Uruguay. *Mar. Pollut. Bull.* 107, 71–76.
565 <https://doi.org/10.1016/j.marpolbul.2016.04.016>

566 Leslie, H.A., Brandsma, S.H., van Velzen, M.J.M., Vethaak, A.D., 2017. Microplastics en route:
567 Field measurements in the Dutch river delta and Amsterdam canals, wastewater treatment
568 plants, North Sea sediments and biota. *Environ. Int.* 101, 133–142.
569 <https://doi.org/10.1016/j.envint.2017.01.018>

570 Lima, A.R.A., Barletta, M., Costa, M.F., 2015. Seasonal distribution and interactions between
571 plankton and microplastics in a tropical estuary. *Estuar. Coast. Shelf Sci.* 165, 213–225.
572 <https://doi.org/10.1016/j.ecss.2015.05.018>

573 Losos, B., Penaz, M., Kubièkova, J., 1980. Food and growth of fishes of the Jihlava river. *Acta*

574 Sc. Nat. 14, 1–46.

575 Mann, R.H.K., 1976. Observations on the age, growth, reproduction and food of the chub
576 *Squalius cephalus* (L.) in the River Stour, Dorset. J. Fish Biol. 8, 265–288.
577 <https://doi.org/10.1111/j.1095-8649.1976.tb03950.x>

578 Mason, S.A., Garneau, D., Sutton, R., Chu, Y., Ehmann, K., Barnes, J., Fink, P., Papazissimos,
579 D., Rogers, D.L., 2016. Microplastic pollution is widely detected in US municipal
580 wastewater treatment plant effluent. Environ. Pollut. 218, 1045–1054.
581 <https://doi.org/10.1016/j.envpol.2016.08.056>

582 Mazurais, D., Ernande, B., Quazuguel, P., Severe, A., Huelvan, C., Madec, L., Mouchel, O.,
583 Soudant, P., Robbens, J., Huvet, A., Zambonino-Infante, J., 2015. Evaluation of the impact
584 of polyethylene microbeads ingestion in European sea bass (*Dicentrarchus labrax*) larvae.
585 Mar. Environ. Res. 112, 78–85. <https://doi.org/10.1016/j.marenvres.2015.09.009>

586 McGoran, A.R., Clark, P.F., Morritt, D., 2017. Presence of microplastic in the digestive tracts of
587 European flounder, *Platichthys flesus*, and European smelt, *Osmerus eperlanus*, from the
588 River Thames. Environ. Pollut. 220, 744–751. <https://doi.org/10.1016/j.envpol.2016.09.078>

589 Michel, P., Oberdoff, T., 1995. Feeding habits of fourteen European freshwater fish species.
590 Cybium 19, 5–46.

591 Morritt, D., Stefanoudis, P. V, Pearce, D., Crimmen, O.A., Clark, P.F., 2014. Plastic in the
592 Thames: A river runs through it. Mar. Pollut. Bull. 78, 196–200.

593 Murphy, F., Russell, M., Ewins, C., Quinn, B., 2017. The uptake of macroplastic & microplastic
594 by demersal & pelagic fish in the Northeast Atlantic around Scotland. Mar. Pollut. Bull. 122,
595 353–359. <https://doi.org/10.1016/j.marpolbul.2017.06.073>

596 Nastova-Grogjioska, R., Kostov, V., Georgiev, S., 1997. Nutrition of chub *Leuciscus cephalus*
597 (Linnaeus, 1758) from river Babuna. Ribarstvo 55, 53–65.

598 Ory, N.C., Sobral, P., Ferreira, J.L., Thiel, M., 2017. Amberstripe scad *Decapterus muroadsi*
599 (Carangidae) fish ingest blue microplastics resembling their copepod prey along the coast of
600 Rapa Nui (Easter Island) in the South Pacific subtropical gyre. *Sci. Total Environ.* 586, 430–
601 437. <https://doi.org/10.1016/j.scitotenv.2017.01.175>

602 Pacini, N., Abate, V., Brambilla, G., De Felip, E., De Filippis, S.P., De Luca, S., di Domenico,
603 A., D’Orsi, A., Forte, T., Fulgenzi, A.R., Iacovella, N., Luiselli, L., Miniero, R., Iamiceli,
604 A.L., 2013. Polychlorinated dibenzodioxins, dibenzofurans, and biphenyls in fresh water
605 fish from Campania Region, southern Italy. *Chemosphere* 90, 80–88.
606 <https://doi.org/10.1016/j.chemosphere.2012.07.039>

607 Pazos, R.S., Maiztegui, T., Colautti, D.C., Paracampo, A.H., Gómez, N., 2017. Microplastics in
608 gut contents of coastal freshwater fish from Río de la Plata estuary. *Mar. Pollut. Bull.* 122,
609 85–90. <https://doi.org/10.1016/j.marpolbul.2017.06.007>

610 Phillips, M.B., Bonner, T.H., 2015. Occurrence and amount of microplastic ingested by fishes in
611 watersheds of the Gulf of Mexico. *Mar. Pollut. Bull.*
612 <https://doi.org/10.1016/j.marpolbul.2015.08.041>

613 Pierce, K.E., Harris, R.J., Larned, L.S., Pokras, M.A., 2004. Obstruction and starvation associated
614 with plastic ingestion in a Northern Gannet *Morus bassanus* and a Greater Shearwater
615 *Puffinus gravis*. *Mar. Ornithol.* 32, 187–189.

616 Pirc, U., Vidmar, M., Mozer, A., Kržan, A., 2016. Emissions of microplastic fibers from
617 microfiber fleece during domestic washing. *Environ. Sci. Pollut. Res.* 23, 22206–22211.
618 <https://doi.org/10.1007/s11356-016-7703-0>

619 Piria, M., Treer, T., Aničić, I., Safner, R., Odak, T., 2005. The natural diet of five cyprinid fish
620 species. *Agric. Conspec. Sci.* 70, 21–28.

621 Rech, S., Macaya-Caquilpán, V., Pantoja, J.F., Rivadeneira, M.M., Jofre Madariaga, D., Thiel,

622 M., 2014. Rivers as a source of marine litter - A study from the SE Pacific. Mar. Pollut.
623 Bull. 82, 66–75.

624 Rochman, C.M., Hoh, E., Hentschel, B.T., Kaye, S., 2013. Long-term field measurement of
625 sorption of organic contaminants to five types of plastic pellets: Implications for plastic
626 marine debris. Environ. Sci. Technol. 47, 1646–1654.

627 Rombout, J.H.W.M., Taverne-Thiele, A.J., Villena, M.I., 1993. The gut-associated lymphoid
628 tissue (GALT) of carp (*Cyprinus carpio* L.): An immunocytochemical analysis. Dev. Comp.
629 Immunol. 17, 55–66. [https://doi.org/10.1016/0145-305X\(93\)90015-I](https://doi.org/10.1016/0145-305X(93)90015-I)

630 Ryan, P.G., 1989. The effects of ingested plastic and other marine debris on seabirds, in:
631 Shomura, R.S., Godfrey, M.L. (Eds.), Proceedings of the Second International Conference
632 on Marine Debris. Honolulu, Hawaii, pp. 623–634.

633 Sanchez, W., Bender, C., Porcher, J.-M., 2014. Wild gudgeons (*Gobio gobio*) from French rivers
634 are contaminated by microplastics: preliminary study and first evidence. Environ. Res. 128,
635 98–100. <https://doi.org/10.1016/j.envres.2013.11.004>

636 Setälä, O., Fleming-Lehtinen, V., Lehtiniemi, M., 2014. Ingestion and transfer of microplastics in
637 the planktonic food web. Environ. Pollut. 185, 77–83.
638 <https://doi.org/10.1016/j.envpol.2013.10.013>

639 Silva-Cavalcanti, J.S., Silva, J.D.B., França, E.J. de, Araújo, M.C.B. de, Gusmão, F., 2017.
640 Microplastics ingestion by a common tropical freshwater fishing resource. Environ. Pollut.
641 221, 218–226. <https://doi.org/10.1016/j.envpol.2016.11.068>

642 Steffens, K.-J., 1995. Persorption — Criticism and Agreement as Based upon In Vitro and In
643 Vivo Studies on Mammals, in: Absorption of Orally Administered Enzymes. Springer Berlin
644 Heidelberg, Berlin, Heidelberg, pp. 9–21. https://doi.org/10.1007/978-3-642-79511-4_2

645 Teuten, E.L., Saquing, J.M., Knappe, D.R.U., Barlaz, M.A., Jonsson, S., Bjorn, A., Rowland,

646 S.J., Thompson, R.C., Galloway, T.S., Yamashita, R., Ochi, D., Watanuki, Y., Moore, C.,
647 Viet, P.H., Tana, T.S., Prudente, M., Boonyatumanond, R., Zakaria, M.P., Akkhavong, K.,
648 Ogata, Y., Hirai, H., Iwasa, S., Mizukawa, K., Hagino, Y., Imamura, A., Saha, M., Takada,
649 H., 2009. Transport and release of chemicals from plastics to the environment and to
650 wildlife. *Philos. Trans. R. Soc. B Biol. Sci.* 364, 2027–2045.
651 <https://doi.org/10.1098/rstb.2008.0284>

652 Van Cauwenberghe, L., Janssen, C.R., 2014. Microplastics in bivalves cultured for human
653 consumption. *Environ. Pollut.* 193, 65–70. <https://doi.org/10.1016/j.envpol.2014.06.010>

654 Vendel, A.L., Bessa, F., Alves, V.E.N., Amorim, A.L.A., Patrício, J., Palma, A.R.T., 2017.
655 Widespread microplastic ingestion by fish assemblages in tropical estuaries subjected to
656 anthropogenic pressures. *Mar. Pollut. Bull.* 117, 448–455.
657 <https://doi.org/10.1016/j.marpolbul.2017.01.081>

658 Volkheimer, G., 1975. Hematogenous dissemination of ingested polyvinyl chloride particles.
659 *Ann. N. Y. Acad. Sci.* 246, 164–71.

660 Von Moos, N., Burkhardt-Holm, P., Köhler, A., 2012. Uptake and effects of microplastics on
661 cells and tissue of the blue mussel *Mytilus edulis* L. after an experimental exposure. *Environ.*
662 *Sci. Technol.* 46, 11327–11335.

663 Wang, W., Ndungu, A.W., Li, Z., Wang, J., 2017. Microplastics pollution in inland freshwaters
664 of China: A case study in urban surface waters of Wuhan, China. *Sci. Total Environ.* 575,
665 1369–1374. <https://doi.org/10.1016/j.scitotenv.2016.09.213>

666 Ward, J.E., Kach, D.J., 2009. Marine aggregates facilitate ingestion of nanoparticles by
667 suspension-feeding bivalves. *Mar. Environ. Res.* 68, 137–142.
668 <https://doi.org/10.1016/j.marenvres.2009.05.002>

669 Watts, A.J.R., Lewis, C., Goodhead, R.M., Beckett, S.J., Moger, J., Tyler, C.R., Galloway, T.S.,

670 2014. Uptake and retention of microplastics by the shore crab *Carcinus maenas*. Environ.
671 Sci. Technol. 48, 8823–8830.

672 Woodall, L.C., Sanchez-Vidal, A., Canals, M., Paterson, G.L.J., Coppock, R., Sleight, V.,
673 Calafat, A., Rogers, A.D., Narayanaswamy, B.E., Thompson, R.C., 2014. The deep sea is a
674 major sink for microplastic debris. R. Soc. Open Sci. 1, 140317–140317.
675 <https://doi.org/10.1098/rsos.140317>

676 Wright, S.L., Kelly, F.J., 2017. Plastic and Human Health: A Micro Issue? Environ. Sci. Technol.
677 51, 6634–6647. <https://doi.org/10.1021/acs.est.7b00423>

678

679